

# Appendix A: Phosphorus Index Worksheet (Complete for each field or crop)

Site/Field: <u>North Dicot</u> <u>Corn</u> <u>2019</u>								
Site Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0, 1, 2, 4, 8)	Weight Factor	Weighted Risk
Soil Erosion	N/A	<5 tons/ac/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	>15 tons/ac/yr	1	X 1.5	1.5
Furrow Irrigation Erosion	N/A	Tailwater recovery, QS>6 very erodible soils, or QS>10 other soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QA>6 for very erodible soils	0	X 1.5	0
Sprinkler Irrigation Erosion	All sites 0-3% slope, all sandy sites, or site evaluation indicates little or no runoff, large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8%, large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	X 0.5	0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1	X 0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	1	X 1.0	<u>1</u>
Commercial P Fertilizer Application Method	None Applied	Placed with planter or injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during the growing season	Incorporated >3 months before crop or surface applied <3 months before crop emerges	Surface applied >3 months before crop emerges	2	X 1.0	2
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	1	X 1.0	1
Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season.	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges.	2	X 1.0	<u>2</u>
Organic P Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	4	X 1.0	<u>4</u>
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or applications are directly into concentrated surface water flow areas.	1	X 1.0	<u>1</u>
Total Phosphorus Index Value: <u>13</u>								

# Appendix A: Phosphorus Index Worksheet (Complete for each field or crop)

Site/Field: <u>South</u> <u>Corn</u> <u>2014</u>								
Site Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0, 1, 2, 4, 8)	Weight Factor	Weighted Risk
Soil Erosion	N/A	<5 tons/ac/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	>15 tons/ac/yr	1	X 1.5	1.5
Furrow Irrigation Erosion	N/A	Tailwater recovery, QS>6 very erodible soils, or QS>10 other soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QA>6 for very erodible soils	0	X 1.5	0
Sprinkler Irrigation Erosion	All sites 0-3% slope, all sandy sites, or site evaluation indicates little or no runoff, large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8%, large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	X 0.5	0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1	X 0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	1	X 1.0	<del>0.5</del> /
Commercial P Fertilizer Application Method	None Applied	Placed with planter or injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during the growing season	Incorporated >3 months before crop or surface applied <3 months before crop emerges	Surface applied >3 months before crop emerges	2	X 1.0	2
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	1	X 1.0	1
Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season.	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges.	2	X 1.0	2
Organic P Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	4	X 1.0	4
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or applications are directly into concentrated surface water flow areas.	1	X 1.0	1
Total Phosphorus Index Value: <u>13</u>								



# Appendix A: Phosphorus Index Worksheet (Complete for each field or crop)

Site/Field: <u>West</u> <u>Alfalfa</u> <u>2014</u>								
Site Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0, 1, 2, 4, 8)	Weight Factor	Weighted Risk
Soil Erosion	N/A	<5 tons/ac/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	>15 tons/ac/yr	1	X 1.5	1.5
Furrow Irrigation Erosion	N/A	Tailwater recovery, QS>6 very erodible soils, or QS>10 other soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QA>6 for very erodible soils	0	X 1.5	0
Sprinkler Irrigation Erosion	All sites 0-3% slope, all sandy sites, or site evaluation indicates little or no runoff, large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8%, large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	X 0.5	0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1	X 0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	1	X 1.0	1
Commercial P Fertilizer Application Method	None Applied	Placed with planter or injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during the growing season	Incorporated >3 months before crop or surface applied <3 months before crop emerges	Surface applied >3 months before crop emerges	0	X 1.0	0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	1	X 1.0	1
Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season.	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges.	2	X 1.0	2
Organic P Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	4	X 1.0	4
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or applications are directly into concentrated surface water flow areas.	1	X 1.0	1
Total Phosphorus Index Value: <u>11</u>								

**Appendix A: Phosphorus Index Worksheet (Complete for each field or crop)**

Site/Field: <u>East Pivot</u> <u>Alfalfa</u> <u>2019</u>								
Site Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0, 1, 2, 4, 8)	Weight Factor	Weighted Risk
Soil Erosion	N/A	<5 tons/ac/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	>15 tons/ac/yr	1	X 1.5	1.5
Furrow Irrigation Erosion	N/A	Tailwater recovery, QS>6 very erodible soils, or QS>10 other soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QA>6 for very erodible soils	0	X 1.5	0
Sprinkler Irrigation Erosion	All sites 0-3% slope, all sandy sites, or site evaluation indicates little or no runoff, large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8%, large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	X 0.5	0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1	X 0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	1	X 1.0	1
Commercial P Fertilizer Application Method	None Applied	Placed with planter or injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during the growing season	Incorporated >3 months before crop or surface applied <3 months before crop emerges	Surface applied >3 months before crop emerges	0	X 1.0	0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	1	X 1.0	1
Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season.	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges.	2	X 1.0	2
Organic P Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	4	X 1.0	4
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or applications are directly into concentrated surface water flow areas.	1	X 1.0	1
Total Phosphorus Index Value: <u>11</u>								



# Appendix A: Phosphorus Index Worksheet (Complete for each field or crop)

Site/Field: <u>Bower Alfalfa 2019</u>								
Site Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0, 1, 2, 4, 8)	Weight Factor	Weighted Risk
Soil Erosion	N/A	<5 tons/ac/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	>15 tons/ac/yr	1	X 1.5	1.5
Furrow Irrigation Erosion	N/A	Tailwater recovery, QS>6 very erodible soils, or QS>10 other soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QA>6 for very erodible soils	0	X 1.5	0
Sprinkler Irrigation Erosion	All sites 0-3% slope, all sandy sites, or site evaluation indicates little or no runoff, large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8%, large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	X 0.5	0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1	X 0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	1	X 1.0	1
Commercial P Fertilizer Application Method	None Applied	Placed with planter or injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during the growing season	Incorporated >3 months before crop or surface applied <3 months before crop emerges	Surface applied >3 months before crop emerges	0	X 1.0	0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	1	X 1.0	1
Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season.	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges.	2	X 1.0	2
Organic P Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	4	X 1.0	4
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or applications are directly into concentrated surface water flow areas.	1	X 1.0	1
Total Phosphorus Index Value: <u>11</u>								

# Appendix A: Phosphorus Index Worksheet (Complete for each field or crop)

Site/Field: <u>South Outground</u> <u>A/F/Grass</u> <u>2014</u>								
Site Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0, 1, 2, 4, 8)	Weight Factor	Weighted Risk
Soil Erosion	N/A	<5 tons/ac/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	>15 tons/ac/yr	1	X 1.5	1.5
Furrow Irrigation Erosion	N/A	Tailwater recovery, QS>6 very erodible soils, or QS>10 other soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QA>6 for very erodible soils	2	X 1.5	3
Sprinkler Irrigation Erosion	All sites 0-3% slope, all sandy sites, or site evaluation indicates little or no runoff, large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8%, large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	X 0.5	0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1	X 0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	1	X 1.0	1
Commercial P Fertilizer Application Method	None Applied	Placed with planter or injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during the growing season	Incorporated >3 months before crop or surface applied <3 months before crop emerges	Surface applied >3 months before crop emerges	0	X 1.0	0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	1	X 1.0	1
Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season.	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges.	2	X 1.0	2
Organic P Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	4	X 1.0	4
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or applications are directly into concentrated surface water flow areas.	1	X 1.0	1
Total Phosphorus Index Value: <u>14</u>								



**Appendix A: Phosphorus Index Worksheet (Complete for each field or crop)**

Site/Field: <u>Bower Outground</u> <u>RIF/Grass</u> <u>2019</u>								
Site Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0, 1, 2, 4, 8)	Weight Factor	Weighted Risk
Soil Erosion	N/A	<5 tons/ac/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	>15 tons/ac/yr	1	X 1.5	1.5
Furrow Irrigation Erosion	N/A	Tailwater recovery, QS>6 very erodible soils, or QS>10 other soils	QS>10 for erosion resistant soils	QS>10 for erodible soils	QA>6 for very erodible soils	2	X 1.5	3
Sprinkler Irrigation Erosion	All sites 0-3% slope, all sandy sites, or site evaluation indicates little or no runoff, large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8%, large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	X 0.5	0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1	X 0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	1	X 1.0	1
Commercial P Fertilizer Application Method	None Applied	Placed with planter or injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during the growing season	Incorporated >3 months before crop or surface applied <3 months before crop emerges	Surface applied >3 months before crop emerges	0	X 1.0	0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	1	X 1.0	1
Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season.	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges.	2	X 1.0	2
Organic P Application Rate	None Applied	<30 lbs/ac P2O5	31-90 lbs/ac P2O5	91-150 lbs/ac P2O5	>150 lbs/ac P2O5	4	X 1.0	4
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or applications are directly into concentrated surface water flow areas.	1	X 1.0	1
Total Phosphorus Index Value: <u>14</u>								